

REFERENCES

1. For selected reviews on cooperative catalysis, see: (a) Ma, J. A.; Cahard, D. Towards perfect catalytic asymmetric synthesis: dual activation of the electrophile and the nucleophile. *Angew. Chem., Int. Ed.* **2004**, *43*, 4566-4583. (b) Kanai, M.; Kato, N.; Ichikawa, E.; Shibasaki, M. Power of cooperativity: Lewis acid-Lewis base bifunctional asymmetric catalysis. *Synlett* **2005**, 1491-1508. (c) Yamamoto, H.; Futatsugi, K. Designer acids: combined acid catalysis for asymmetric synthesis. *Angew. Chem., Int. Ed.* **2005**, *44*, 1924-1942. (d) Akiyama, T. Stronger Brønsted acids. *Chem. Rev.* **2007**, *107*, 5744-5758. (e) Terada, M. Binaphthol-derived phosphoric acid as a versatile catalyst for enantioselective carbon-carbon bond forming reactions. *Chem. Commun.* **2008**, 4097-4112. (f) Paull, D. H.; Abraham, C. J.; Scerba, M. T.; Alden-Danforth, E.; Lectka, T. Bifunctional asymmetric catalysis: cooperative Lewis acid/base systems. *Acc. Chem. Res.* **2008**, *41*, 655-663. (g) Kobayashi, S.; Mori, Y.; Fossey, J. S.; Salter, M. M. Catalytic enantioselective formation of C-C bonds by addition to imines and hydrazones: a ten-year update. *Chem. Rev.* **2011**, *111*, 2626-2704. (h) Fulton, J. R.; Glover, J. E.; Kamara, L.; Rowlands, G. J. Facile synthesis of planar chiral *N*-oxides and their use in Lewis base catalysis. *Chem. Commun.* **2011**, *47*, 433-435. (i) Allen, A. E.; MacMillan, D. W. C. Synergistic catalysis: a powerful synthetic strategy for new reaction development. *Chem. Sci.* **2012**, *3*, 633-658. (j) Park, J.; Hong, S. Cooperative bimetallic catalysis in asymmetric transformations. *Chem. Soc. Rev.* **2012**, *41*, 6931-6943. (k) Cohen, D. T.; Scheidt, K. A. Cooperative Lewis acid/*N*-heterocyclic carbene catalysis. *Chem. Sci.* **2012**, *3*, 53-57. (l) Inamdar, S. M.; Shinde, V. S.; Patil, N. T. Enantioselective cooperative catalysis. *Org. Biomol. Chem.* **2015**, *13*, 8116-8162. (m) Trost, B. M.; Bartlett, M. J. ProPhenol-catalyzed asymmetric additions by spontaneously assembled dinuclear main group metal complexes. *Acc. Chem. Res.* **2015**, *48*, 688-701. (n) Shibasaki, M.; Kumagai, N. in *Cooperative Catalysis: Designing Efficient Catalysts for Synthesis*, Peters, R., Eds.; Wiley-VCH: New York, **2015**; Chapter 1. (o) Skubi, K. L.; Blum, T. R.; Yoon, T. P. Dual catalysis strategies in photochemical synthesis. *Chem. Rev.* **2016**, *116*, 10035-10074. (p) Afewerki, S.; Córdova, A. Combinations of aminocatalysts and metal catalysis: a powerful cooperative approach in selective organic synthesis. *Chem. Rev.* **2016**, *116*, 13512-13570. (q) Kim, D.-S.; Park, W.-J.; Jun, C.-H. Metal-organic cooperative catalysis in C-H and C-C bond activation. *Chem. Rev.* **2017**, *117*, 8977-9015. (r) Romiti, F.; del Pozo, J.; Paioti, P. H. S.; Gonsales, S. A.; Li, X.; Hartrampf, F. W. W.; Hoveyda, A. H. Different Strategies for Designing Dual-Catalytic Enantioselective Processes: From Fully Cooperative to Non-cooperative Systems. *J. Am. Chem. Soc.* **2019**, *141*, 17952-17961.
2. For reviews of hydrogen bonding catalysis, see: (a) Schreiner, P. R. Metal-free organocatalysis through explicit hydrogen bonding interactions. *Chem. Soc. Rev.* **2003**, *32*, 289-296. (b) Takemoto, Y. Recognition and activation by ureas and thioureas: stereoselective reactions using ureas and thioureas as hydrogen-bonding donors. *Org. Biomol. Chem.* **2005**, *3*, 4299-4306. (c) Connon, S. J. Organocatalysis mediated by (thio)urea derivatives. *Chem. Eur. J.* **2006**, *12*, 5418-5427. (d) Taylor, M. S.; Jacobsen, E. N. Asymmetric catalysis by chiral hydrogen-bond donors. *Angew. Chem., Int. Ed.* **2006**, *45*, 1520-1543. (e) Doyle, A. G.; Jacobsen, E. N. Small-molecule H-bond donors in asymmetric catalysis. *Chem. Rev.* **2007**, *107*, 5713-5743. (f) Mukherjee, S.; Yang, J. W.; Hoffmann, S.; List, B. Asymmetric enamine catalysis. *Chem. Rev.* **2007**, *107*, 5471-5569. (g) Knowles, R. R.; Jacobsen, E. N. Attractive noncovalent interactions in asymmetric catalysis: Links between enzymes and small molecule catalysts. *Proc. Natl. Acad. Sci. U. S. A.* **2010**, *107*, 20678-20685.

3. For reviews of cooperative two-center catalysis, see: (a) Karlin, K. D. Metalloenzymes, structural motifs, and inorganic models. *Science* **1993**, *261*, 701-708. (b) Strater, N.; Lipscomb, W. N.; Klabunde, T.; Krebs, B. Two-metal ion catalysis in enzymatic acyl- and phosphoryl-transfer reactions. *Angew. Chem., Int. Ed.* **1996**, *35*, 2024-2055. (c) Wilcox, D. E. Binuclear metallohydrolases. *Chem. Rev.* **1996**, *96*, 2435-2458. (d) Jedrzejewski, M. J.; Setlow, P. Comparison of the binuclear metalloenzymes diphosphoglycerate-independent phosphoglycerate mutase and alkaline phosphatase: Their mechanism of catalysis via a phosphoserine intermediate. *Chem. Rev.* **2001**, *101*, 607-618.
4. For selected reviews on iminium and enamine catalysis, see: (a) Notz, W.; Tanaka, F.; Barbas, C. F., 3rd Enamine-based organocatalysis with proline and diamines: the development of direct catalytic asymmetric Aldol, Mannich, Michael, and Diels-alder reactions. *Acc. Chem. Res.* **2004**, *37*, 580-91. (b) Erkkilä, A.; Majander, I.; Pihko, P. M. Iminium Catalysis. *Chem. Rev.* **2007**, *107*, 5416-5470. (c) Mukherjee, S.; Yang, J. W.; Hoffmann, S.; List, B. Asymmetric Enamine Catalysis. *Chem. Rev.* **2007**, *107*, 5471-5569. (d) Zou, Y.-Q.; Hörmann, F. M.; Bach, T. Iminium and enamine catalysis in enantioselective photochemical reactions. *Chem. Soc. Rev.* **2018**, *47*, 278-290.
5. For representative examples of enantioselective reactions by enamine catalysis, see: (a) Barbas, C. F.; Heine, A.; Zhong, G.; Hoffmann, T.; Gramatikova, S.; Björnstedt, R.; List, B.; Anderson, J.; Stura, E. A.; Wilson, I. A.; Lerner, R. A. Immune Versus Natural Selection: Antibody Aldolases with Enzymic Rates But Broader Scope. *Science* **1997**, *278*, 2085-2092. (b) List, B.; Lerner, R. A.; Barbas, C. F. Proline-Catalyzed Direct Asymmetric Aldol Reactions. *J. Am. Chem. Soc.* **2000**, *122*, 2395-2396. (c) Northrup, A. B.; MacMillan, D. W. The first direct and enantioselective cross-aldol reaction of aldehydes. *J. Am. Chem. Soc.* **2002**, *124*, 6798-9. (d) Kumaragurubaran, N.; Juhl, K.; Zhuang, W.; Bøgevig, A.; Jørgensen, K. A. Direct *L*-Proline-Catalyzed Asymmetric α -Amination of Ketones. *J. Am. Chem. Soc.* **2002**, *124*, 6254-6255. (e) List, B.; Pojarliev, P.; Biller, W. T.; Martin, H. J. The Proline-Catalyzed Direct Asymmetric Three-Component Mannich Reaction: Scope, Optimization, and Application to the Highly Enantioselective Synthesis of 1,2-Amino Alcohol. *J. Am. Chem. Soc.* **2002**, *124*, 827. (f) Córdova, A.; Notz, W.; Zhong, G.; Betancort, J. M.; Barbas, C. F., III. A Highly Enantioselective Amino Acid-Catalyzed Route to Functionalized α -Amino Acids. *J. Am. Chem. Soc.* **2002**, *124*, 1842. (g) Córdova, A.; Watanabe, S.-i.; Tanaka, F.; Notz, W.; Barbas, C. F., III. A Highly Enantioselective Route to Either Enantiomer of Both α - and β -Amino Acid Derivatives. *J. Am. Chem. Soc.* **2002**, *124*, 1866. (h) Kano, T.; Yamaguchi, Y.; Tokuda, O.; Maruoka, K. anti-Selective Direct Asymmetric Mannich Reactions Catalyzed by Axially Chiral Amino Sulfonamide as an Organocatalyst. *J. Am. Chem. Soc.* **2005**, *127*, 16408. (i) Mitsumori, S.; Zhang, H.; Cheong, P. H.-Y.; Houk, K. N.; Tanaka, F.; Barbas, C. F., III. Direct Asymmetric anti-Mannich-Type Reactions Catalyzed by a Designed Amino Acid. *J. Am. Chem. Soc.* **2006**, *128*, 1040. (j) Hayashi, Y.; Urushima, T.; Aratake, S.; Okano, T.; Obi, K. Organic Solvent-Free, Enantio- and Diastereoselective, Direct Mannich Reaction in the Presence of Water. *Org. Lett.* **2008**, *10*, 21. (k) Zhao, Y.; Pan, Y.; Liu, H.; Yang, Y.; Jiang, Z.; Tan, C.-H. Fluorinated Aromatic Ketones as Nucleophiles in the Asymmetric Organocatalytic Formation of C–C and C–N Bonds: A Facile Route to the Construction of Fluorinated Quaternary Stereogenic Centers. *Chem. Eur. J.* **2011**, *17*, 3571.
6. For selected reviews on FLP chemistry, see: (a) *Frustrated Lewis Pairs I: Uncovering and Understanding*; Stephan, D. W.; Erker, G. Eds.; Springer: Berlin, 2013; Vol. 332, p 1-289. (b) *Frustrated Lewis Pairs II: Expanding the Scope*; Erker, G.; Stephan, D. W. Eds.; Springer: Berlin, 2013; Vol. 334, p 1-311. (c) Ashley, A. E.; O'Hare, D. FLP-mediated activations and reductions of CO₂ and CO. *Top. Curr. Chem.* **2013**, *334*, 191-218. (d) Feng, X.; Du, H. Metal-free asymmetric

- hydrogenation and hydrosilylation catalyzed by frustrated Lewis pairs. *Tetrahedron Lett.* **2014**, *55*, 6959-6964. (e) Stephan, D. W.; Erker, G. Frustrated Lewis pair chemistry: Development and perspectives. *Angew. Chem., Int. Ed.* **2015**, *54*, 6400-6441. (f) Stephan, D. W. Frustrated Lewis pairs. *J. Am. Chem. Soc.* **2015**, *137*, 10018-10032. (g) Stephan, D. W. The broadening reach of frustrated Lewis pair chemistry. *Science* **2016**, *354*, aaf7229. (h) Scott, D. J.; Fuchter, M. J.; Ashley, A. E. Designing effective 'frustrated Lewis pair' hydrogenation catalysts. *Chem. Soc. Rev.* **2017**, *46*, 5689-5700. (i) Lam, J.; Szkop, K. M.; Mosafari, E.; Stephan, D. W. FLP catalysis: main group hydrogenations of organic unsaturated substrates. *Chem. Soc. Rev.* **2019**, *48*, 3592-3612. (j) Stephan, D. W. Catalysis, FLPs, and beyond. *Chem* **2020**, *6*, 1520-1526. (k) Carden, J. L.; Dasgupta, A.; Melen, R. L. Halogenated triarylboranes: synthesis, properties and applications in catalysis. *Chem. Soc. Rev.* **2020**, *49*, 1706-1725. (l) Ma, Y.; Lou, S.-J.; Hou, Z. Electron-deficient boron-based catalysts for C-H bond functionalisation. *Chem. Soc. Rev.* **2021**, *50*, 1945-1967.
7. For reviews on late stage functionalization, see: (a) Cernak, T.; Dykstra, K. D.; Tyagarajan, S.; Vachal, P.; Krska, S. W. The medicinal chemist's toolbox for late stage functionalization of drug-like molecules. *Chem. Soc. Rev.* **2016**, *45*, 546-576. (b) Moir, M.; Danon, J. J.; Reekie, T. A.; Kassiou, M. An overview of late-stage functionalization in today's drug discovery. *Expert Opin. Drug Discov.* **2019**, *14*, 1137-1149. (c) Campos, K. R.; Coleman, P. J.; Alvarez, J. C.; Dreher, S. D.; Garbaccio, R. M.; Terrett, N. K.; Tillyer, R. D.; Truppo, M. D.; Parmee, E. R. The importance of synthetic chemistry in the pharmaceutical industry. *Science* **2019**, *363*, eaat0805. (d) Börgel, J.; Ritter, T. Late-Stage Functionalization. *Chem* **2020**, *6*, 1877-1887. (e) Guillemard, L.; Kaplaneris, N.; Ackermann, L.; Johansson, M. J. Late-stage C-H functionalization offers new opportunities in drug discovery. *Nat. Rev. Chem.* **2021**.
8. For selected examples of FLP-mediated H₂ splitting, see: (a) Welch, G. C.; Juan, R. R. S.; Masuda, J. D.; Stephan, D. W. Reversible, Metal-Free Hydrogen Activation. *Science* **2006**, *314*, 1124-1126. (b) Welch, G. C.; Stephan, D. W. Facile Heterolytic Cleavage of Dihydrogen by Phosphines and Boranes. *J. Am. Chem. Soc.* **2007**, *129*, 1880-1881. (c) Spies, P.; Erker, G.; Kehr, G.; Bergander, K.; Fröhlich, R.; Grimme, S.; Stephan, D. W. Rapid intramolecular heterolytic dihydrogen activation by a four-membered heterocyclic phosphane-borane adduct. *Chem. Commun.* **2007**, 5072-5074. (d) Liu, Y.; Du, H. Metal-Free Borane-Catalyzed Highly Stereoselective Hydrogenation of Pyridines. *J. Am. Chem. Soc.* **2013**, *135*, 12968-12971. (e) Mahdi, T.; Stephan, D. W. Enabling Catalytic Ketone Hydrogenation by Frustrated Lewis Pairs. *J. Am. Chem. Soc.* **2014**, *136*, 15809-15812. (f) Scott, D. J.; Fuchter, M. J.; Ashley, A. E. Nonmetal Catalyzed Hydrogenation of Carbonyl Compounds. *J. Am. Chem. Soc.* **2014**, *136*, 15813-15816.
9. For a selected review on FLP-mediated CO₂ activation, see: Ashley, A. E.; O'Hare, D. FLP-mediated activations and reductions of CO₂ and CO. *Top. Curr. Chem.* **2013**, *334*, 191-217.
10. For selected references on organoborane-mediated hydride abstraction from amines, see: (a) Millot, N.; Santini, C. C.; Fenet, B.; Basset, J. M. Formation and characterization of zwitterionic stereoisomers from the reaction of B(C₆F₅)₃ and NEt₂Ph: (E)- and (Z)-[EtPhN⁺=CHCH₂-B⁻(C₆F₅)₃]. *Eur. J. Inorg. Chem.* **2002**, *2002*, 3328-3335. (b) Focante, F.; Mercandelli, P.; Sironi, A.; Resconi, L. Complexes of tris(pentafluorophenyl)boron with nitrogen-containing compounds: Synthesis, reactivity and metallocene activation. *Coord. Chem. Rev.* **2006**, *250*, 170-188. (c) Dureen, M. A.; Brown, C. C.; Stephan, D. W. Addition of enamines or pyrroles and B(C₆F₅)₃ Frustrated Lewis Pairs to alkynes. *Organometallics* **2010**, *29*, 6422-6432. (d) Schwendemann, S.; Fröhlich, R.; Kehr, G.; Erker, G. Intramolecular frustrated N/B Lewis pairs by enamine hydroboration. *Chem. Sci.* **2011**, *2*, 1842-1849. (e) Tian, J.-J.; Zeng, N.-N.; Liu, N.; Tu, X.-S.; Wang, X.-C. Intramolecular cyclizations

- of vinyl-substituted *N,N*-dialkyl arylamines enabled by borane-assisted hydride transfer. *ACS Catal.* **2019**, *9*, 295–300.
11. For selected references on organoborane-catalyzed transformation of amines into enamines, see: (a) Zhang, J.; Park, S.; Chang, S. Catalytic access to bridged sila-*N*-heterocycles from piperidines via cascade sp^3 and sp^2 C–Si bond formation. *J. Am. Chem. Soc.* **2018**, *140*, 13209–13213. (b) Li, R.; Chen, Y.; Jiang, K.; Wang, F.; Lu, C.; Nie, J.; Chen, Z.; Yang, G.; Chen, Y.-C.; Zhao, Y.; Ma, C. $B(C_6F_5)_3$ -Catalyzed redox-neutral β -alkylation of tertiary amines using *p*-quinone methides via borrowing hydrogen. *Chem. Commun.* **2019**, *55*, 1217–1220. (c) Zhang, J.; Chang, S. *cine*-Silylative ring-opening of α -methyl azacycles enabled by the silylium-induced C–N bond cleavage. *J. Am. Chem. Soc.* **2020**, *142*, 12585–12590. (d) Chen, Y.; Wan, H. L.; Huang, Y.; Liu, S.; Wang, F.; Lu, C.; Nie, J.; Chen, Z.; Yang, G.; Ma, C. $B(C_6F_5)_3$ -Catalyzed β -functionalization of pyrrolidines using isatins via borrowing hydrogen: Divergent access to substituted pyrrolidines and pyrroles. *Org. Lett.* **2020**, *22*, 7797–7803.
 12. (a) Maier, A. F. G.; Tussing, S.; Schneider, T.; Flörke, U.; Qu, Z.-W.; Grimme, S.; Paradies, J. Frustrated Lewis pair catalyzed dehydrogenative oxidation of indolines and other heterocycles. *Angew. Chem., Int. Ed.* **2016**, *55*, 12219–12223. (b) Kojima, M.; Kanai, M. Tris(pentafluorophenyl)borane-catalyzed acceptorless dehydrogenation of *N*-heterocycles. *Angew. Chem., Int. Ed.* **2016**, *55*, 12224–12227.
 13. Shang, M.; Chan, J. Z.; Cao, M.; Chang, Y.; Wang, Q.; Cook, B.; Torker, S.; Wasa, M. C–H Functionalization of Amines via Alkene-Derived Nucleophiles through Cooperative Action of Chiral and Achiral Lewis Acid Catalysts: Applications in Enantioselective Synthesis. *J. Am. Chem. Soc.* **2018**, *140*, 10593–10601.
 14. Chan, J. Z.; Yesilcimen, A.; Cao, M.; Zhang, Y.; Zhang, B.; Wasa, M. Direct Conversion of *N*-Alkylamines to *N*-Propargylamines through C–H Activation Promoted by Lewis Acid/Organocopper Catalysis: Application to Late-Stage Functionalization of Bioactive Molecules. *J. Am. Chem. Soc.* **2020**, *142*, 16493–16505.
 15. For selected reviews on C–H activation reactions, see: (a) Dick, A. R.; Sanford, M. S. Transition metal catalyzed oxidative functionalization of carbon-hydrogen bonds. *Tetrahedron* **2006**, *62*, 2439–2463. (b) Davies, H. M.; Manning, J. R. Catalytic C–H functionalization by metal carbenoid and nitrenoid insertion. *Nature* **2008**, *451*, 417–424. (c) Baudoin, O. Transition metal-catalyzed arylation of unactivated $C(sp^3)$ -H bonds. *Chem. Soc. Rev.* **2011**, *40*, 4902–4911. (d) Haibach, M. C.; Seidel, D. C–H Bond functionalization through intramolecular hydride transfer. *Angew. Chem., Int. Ed.* **2014**, *53*, 5010–5036. (e) Girard, S. A.; Knauber, T.; Li, C.-J. The cross-dehydrogenative coupling of C_{sp^3} -H bonds: A versatile strategy for C–C bond formations. *Angew. Chem., Int. Ed.* **2014**, *53*, 74–100. (f) Daugulis, O.; Roane, J.; Tran, L. D. Bidentate, monoanionic auxiliary-directed functionalization of carbon-hydrogen bonds. *Acc. Chem. Res.* **2015**, *48*, 1053–1064. (g) He, J.; Wasa, M.; Chan, K. S. L.; Shao, Q.; Yu, J.-Q. Palladium-catalyzed transformations of alkyl C–H bonds. *Chem. Rev.* **2017**, *117*, 8754–8786. (h) Chu, J. C. K.; Rovis, T. Complementary strategies for directed $C(sp^3)$ -H functionalization: A comparison of transition-metal-catalyzed activation, hydrogen atom transfer, and carbene/nitrene transfer. *Angew. Chem., Int. Ed.* **2018**, *57*, 62–101. (i) Davies, H. M.; Liao, K. Dirhodium tetracarboxylates as catalysts for selective intermolecular C–H functionalization. *Nat. Rev. Chem.* **2019**, *3*, 347–360. (j) Campos, K. R. Direct sp^3 C–H bond activation adjacent to nitrogen in heterocycles. *Chem. Soc. Rev.* **2007**, *36*, 1069–1084. (k) Mitchell, E. A.; Peschiulli, A.; Lefevre, N.; Meerpoel, L.; Maes, B. U. W. Direct α -functionalization of saturated cyclic amines. *Chem. Eur. J.* **2012**, *18*, 10092–10142. (l) He, C.; Whitehurst, W. G.; Gaunt, M. J.

- Palladium-catalyzed C(sp³)-H bond functionalization of aliphatic amines. *Chem* **2019**, *5*, 1031-1058. (m) Trowbridge, A.; Walton, S. M.; Gaunt, M. J. New strategies for the transition-metal catalyzed synthesis of aliphatic amines. *Chem. Rev.* **2020**, *120*, 2613-2692.
16. Chang, Y.; Cao, M.; Chan, J. Z.; Zhao, C.; Wang, Y.; Yang, R.; Wasa, M. Enantioselective Synthesis of *N*-Alkylamines through β -Amino C-H Functionalization Promoted by Cooperative Actions of B(C₆F₅)₃ and a Chiral Lewis Acid Co-Catalyst. *J. Am. Chem. Soc.* **2021**, *143*, 2441-2455.
17. Chang, Y.; Yesilcimen, A.; Cao, M.; Zhang, Y.; Zhang, B.; Chan, J. Z.; Wasa, M. Catalytic Deuterium Incorporation within Metabolically Stable β -Amino C-H Bonds of Drug Molecules. *J. Am. Chem. Soc.* **2019**, *141*, 14570-14575.